

Public Health Scotland COVID-19 & Winter Statistical Report

As at 14 February 2022

Publication date: 16 February 2022

Contents

Introduction	4
Main Points	6
Incidence of Variants of Concern and Variants Under Investigation	7
COVID-19 Daily Data	8
Lateral Flow Device (LFD) Testing	11
COVID-19 Hospital Admissions	13
Test and Protect	17
COVID-19 Vaccine	25
COVID-19 Vaccination Uptake	26
COVID-19 Vaccine Wastage	27
COVID-19 Vaccine Certification	28
COVID-19 cases, acute hospitalisations, and deaths by vaccine status	29
Hospital/ Wider System Pressures	49
Unscheduled Care	49
Waiting Times	52
Delayed Discharges	53
Wider Impact of COVID-19	54
Contact	55
Further Information	55
Open Data	55
Rate this publication	55
Early access details	55
Appendices	56
Appendix 1: Background information	56
Appendix 2: World Health Organisation (WHO): Contact tracing in the context of COVID-19	56
Appendix 3: Hospital Admissions Notes	57
Appendix 4: Contact Tracing	57
Appendix 5: Lateral Flow Device Testing	58
Appendix 6: Data sources and Limitations	59

This is a Management Information publication

Published management information are non-official statistics. They may not comply with the UK Statistics Authority's Code of Practice with regard to high data quality or high public value but there is a public interest or a specific interest by a specialist user group in accessing these statistics, as there are no associated official statistics available. Users should therefore be aware of the aspects of data quality and caveats surrounding these data, all of which are listed in this document. Therefore, the data presented are subject to change.

Introduction

Since the start of the Coronavirus-19 (COVID-19) outbreak Public Health Scotland (PHS) has been working closely with Scottish Government and health and care colleagues in supporting the surveillance and monitoring of COVID-19 amongst the population. As part of our continuous review of reporting, as of 08 December 2021 Public Health Scotland has implemented changes to the COVID-19 Weekly Report to support the reader in drawing insights from a wider range of existing metrics around COVID-19 and winter pressures.

Caution should be used when making comparisons between metrics; each metric is calculated independently and may cover different time periods or cohorts of the population.

The consolidated report will include the following content weekly:

COVID-19

- Summary of tests and cases
- Contact Tracing
- Hospital and ICU admissions
- Testing in care homes
- COVID-19 vaccination status cases, hospitalisations and deaths
- Covid-19 vaccination uptake summary
- Adhoc reporting on topics such as: Covid-19 and Vaccination in pregnancy, Equality reporting etc.

Hospital/ Wider System Pressures

- Unscheduled Care
- Waiting Times
- Delayed Discharges

Additional charts for a number of variables related to COVID-19 service use in the NHS, including some metrics previously presented in the weekly COVID-19 report, are available to view in our [interactive dashboard](#). These include breakdowns by age, sex and deprivation. The variables currently available on the dashboard include:

- Positive cases per day and cumulative total
- COVID-19 hospital admissions
- COVID-19 patients admitted to ICU admissions
- COVID-19 related contacts to NHS24 and the Coronavirus Helpline
- Community Hubs and Assessment Centres
- Scottish Ambulance Service incidents
- Contact tracing
- Health care workers
- Care homes
- Targeted community testing
- Travel outside of Scotland
- Quarantine Statistics
- NHS Protect Scotland App
- Lateral Flow Device (LFD) Testing

The Public Health Scotland [COVID-19 Daily Dashboard](#) publishes daily updates on the number of positive cases of COVID-19 in Scotland, with charts showing the trend since the start of the outbreak. From 26 February 2021, the Daily Dashboard also includes daily updates on vaccinations for COVID-19 in Scotland.

There is a large amount of data being regularly published regarding COVID-19 (for example, [Coronavirus in Scotland – Scottish Government](#) and [Deaths involving coronavirus in Scotland – National Records of Scotland](#)). This report complements the range of existing data currently available.

In addition, the [COVID-19 State of the Epidemic Report – Scottish Government](#) brings together different sources of evidence and data about the epidemic in Scotland, and shows how Scotland currently compares to the rest of the UK.

Main Points

- In the week ending 13 February 2022, there were 40,873 COVID-19 cases (identified by PCR or LFD), a decrease of 8.1% from the previous week
- In the week ending 06 February 2022, 48,445 individuals were recorded in the contact tracing software, from which 50,245 unique contacts have been traced
- In the week ending 08 February 2022, there were 607 admissions to hospital with a positive COVID-19 test (PCR or LFD), a decrease of 5% from the previous week. The highest number of new admissions are now in those aged 80+
- In the week ending 13 February 2022 there were 21 new admissions to Intensive Care Units (ICUs) with a laboratory confirmed test of COVID-19. This is a decrease of 16.0% from the week ending 06 February 2022
- In the most recent four weeks, from 15 January 2022 to 11 February 2022, the age-standardised rate of hospital admissions per 100,000 were 3.0 to 3.7 times lower in individuals with their third dose or booster dose of vaccine compared to unvaccinated individuals or have received one or two doses of a COVID-19 vaccine

Incidence of Variants of Concern and Variants Under Investigation

The Omicron variant was originally detected in South Africa and now represents the dominant variant in Scotland.

Further information on previous Omicron reporting can be found [here](#).

Public Health Scotland (PHS) continues to monitor COVID-19 Variants of Concern, in collaboration with other Public Health Agencies in the UK.

The latest [information on the number of such variants detected by genomic analyses across the UK](#) is published by UK Health Security Agency (UKHSA).

COVID-19 Daily Data

The Public Health Scotland [COVID-19 Daily Dashboard](#) publishes daily updates (5-days per week, Monday to Friday) on the number of positive cases of COVID-19 in Scotland, with charts showing the trend since the start of the outbreak.


The total number of people within Scotland who have, or have had COVID-19, since the coronavirus outbreak began is unknown. The number of laboratory confirmed cases is likely to be an underestimate of the total number who have, or have had, COVID-19. A person can have multiple tests but will only ever be counted once. The drop in the number of confirmed cases at weekends likely reflects that laboratories are doing fewer tests at the weekend.

On 05 January 2022, the Scottish Government announced that asymptomatic people who return a positive lateral flow test (LFD) no longer have to confirm their positive result with a PCR test.

The statistics presented in this section reflect the change in case definition (an individual's first positive PCR or LFD from 05 January 22). More information available on the Public Health Scotland website [here](#).


- In the week ending 13 February 2022 there were 40,873 COVID-19 cases by PCR or LFD, with a seven-day rolling average of 5,839 cases¹. This is a decrease of 8.1% from the previous week.

Figure 1: Number of positive cases per day with 7-day average



1. Correct as at 14 February, may differ from more recently published data in the previous week's report and on the [COVID-19 Daily Dashboard](#).


Figure 2: Number of positive cases by test type per day, 06 January 2022 to 13 February 2022*



* A positive case is defined as an individual's first positive PCR or first positive LFD test. LFD tests are included in the case definition from 05 January 2022 to reflect the revised testing strategy. LFD positive cases that are followed by a negative PCR result within 48 hours will be excluded. Cases will only be counted once.

Figure 3 below shows the proportion of COVID-19 cases by PCR or LFD, by age group for the most recent three weeks.

Figure 3: Proportion of COVID-19 cases (PCR or LFD) by age group, weeks ending 30 January 2022 – 13 February 2022




The [daily dashboard](#) also includes data on Hospital Admissions (PCR or LFD) and ICU admissions (PCR only) for patients with COVID-19:

- In the week ending 08 February 2022, there were 607 admissions to hospital with a PCR or LFD positive test of COVID-19.
- In the week ending 01 February 2022 there were 21 new admissions to Intensive Care Units (ICUs) for PCR laboratory confirmed COVID-19 patients.

The number of daily positive COVID-19 cases (PCR or LFD) increased from 7,477 to 7,605 between 01 February 2022 and 08 February 2022. During this same time period, the daily positive COVID-19 PCR or LFD hospital admissions has decreased from 105 to 90 (seven-day rolling average). The seven-day average of inpatients in hospital has decreased by 18.0% (from 1,229 to 1,030).

Figure 4: Number of Positive Cases, Admissions and Inpatients, as at 08 February 2022²



2. Please refer to [Appendix 3 - Hospital Admissions Notes](#) for definitions of hospital admissions and inpatients. Note that prior to 6 Jan 2022 cases are PCR only.

Additional charts and data are available to view in the [interactive dashboard](#) accompanying this report.

Data is also monitored and published daily on the [Scottish Government Coronavirus website](#).

Lateral Flow Device (LFD) Testing

Across Scotland, there are numerous testing pathways being rolled out using Lateral Flow Devices (LFD) - a clinically validated swab antigen test taken that does not require a laboratory for processing. This test can produce rapid results within 45 minutes at the location of the test.

Some of the areas using LFD tests are schools, health and social care workers, care homes and more. Public Health Scotland has collected the information on the number of LFD tests carried out across Scotland and will now publish this information weekly. This section is the totality of LFD across Scotland and across strategies.

On 05 January 2022, the Scottish Government announced that people who do not have symptoms would no longer be asked to take a polymerase chain reaction (PCR) test to confirm a positive Lateral Flow Device (LFD) result. Instead, anyone with a positive LFD, who does not have symptoms, should report the result online as soon as the test is done. In order to ensure that we continue to provide the most accurate information, changes have been made to the national COVID-19 case definition to reflect this revised testing strategy.

Since 19 November 2020, there have been 23,121,318 LFD tests carried out in Scotland, of which 380,860 were positive (1.6%). Figure 5 below shows the weekly trend of tests carried out from week ending 29 November 2020 to 13 February 2022.

There has been a 5.3% decrease in the number of tests carried out from the week ending 06 February 2022 to the week ending 13 February 2022. Table 1 shows the number of LFD tests carried out in Scotland by testing group.

More detailed information can be found within the LFD section on our [interactive dashboard](#). For additional details on Lateral Flow Device Tests, please see - [Appendix 5 – Lateral Flow Device Testing](#)

Figure 5: Trend of LFD tests carried out in Scotland from 29 November 2020 to 13 February 2022




Table 1: Number of LFD¹⁰ tests by Test Group 19 November 2020 – 13 February 2022

Test Group	Test Reason	Number of tests	Number of positive tests	% LFT positive
Care Home Testing	Care Home - Visiting Professional	68,327	246	0.4%
	Care Home - Visitor	937,442	1,465	0.2%
	Care Home Staff	2,157,552	5,874	0.3%
Community Testing	Community Testing	107,912	993	0.9%
Education Testing	Combined School Staff	64,779	374	0.6%
	ELC Staff	378,001	4,027	1.1%
	Primary School Staff	1,752,927	13,633	0.8%
	Secondary School Pupils	1,112,554	18,531	1.7%
	Secondary School Staff	1,002,132	6,578	0.7%
	University Staff	16,363	305	1.9%
	University Students	82,401	2,404	2.9%
Healthcare Testing	University Testing Site	97,560	387	0.4%
	Healthcare Worker	3,430,964	18,340	0.5%
Social Care Testing	Primary Care And Independent Contractors	276,858	1,463	0.5%
	Children, Young People And Mental Health	1,060	0	0%
	NSS Portal Social Care	868,279	4,138	0.5%
	Residential Homes	18,094	95	0.5%
Universal Offer	Support Services	54,052	886	1.6%
	Attend An Event	1,150,669	5,679	0.5%
	High Cases In Local Area	828,990	25,566	3.1%
	Lives With Someone Who Is Shielding	107,152	3,147	2.9%
	Travel Within UK	291,761	2,424	0.8%
Workplace Testing	Universal Offer	4,512,316	188,323	4.2%
	Private Sector	33,292	223	0.7%
	Public Sector	87,270	531	0.6%
	Quarantine Hotel Staff/Security Personnel	5,999	263	4.4%
	Third Sector	6,309	37	0.6%
Other	UK Gov Other	2,747,148	46,699	1.7%
	Other	923,155	28,229	3.1%
Total	Total	23,121,318	380,860	1.6%

Data extracted: 14 February

Please note some of the data is suppressed due to disclosure methodology being applied to protect staff confidentiality.

COVID-19 Hospital Admissions

Hospital Admissions 'with' COVID-19

Since the start of the pandemic Public Health Scotland have been reporting on the number of people in acute hospitals with recently confirmed COVID-19. These admissions are identified from Rapid and Preliminary Inpatient Data (RAPID) and defined as the following: A patient's first positive PCR or LFD test (LFD from the 5th of January) for COVID-19 up to 14 days prior to admission to hospital, on the day of their admission or during their stay in hospital. If a patient's first positive test is after their date of discharge from hospital, they are not included in the analysis.

The statistics presented in this section reflect the change in case definition (an individual's first positive PCR or LFD from 05 January 22). More information available on the Public Health Scotland website [here](#).

It is important to note, that the figures presented below may include patients being admitted and treated in hospital for reasons other than COVID-19. Supplementary analysis on COVID-19 related acute hospital admissions by vaccine status is also available within the [COVID-19 cases, acute hospitalisations, and deaths by vaccine status](#) section of this report.

Figure 6 below shows the weekly trend of hospital admissions with COVID-19 from week ending 05 January 2021 to 08 February 2022.

Figure 6: Trend of hospital admissions 'with' COVID-19 in Scotland




Table 2 below shows a breakdown of people admitted to hospital across all ages and by age group for the most recent four weeks. Data from March 2021 is available on the [Covid Statistical Report website](#).

The statistics presented in this section reflect the change in case definition (an individual's first positive PCR of LFD from 05 January 22). More information available on the Public Health Scotland website [here](#).

Table 2: COVID-19 hospital admissions by age as at 08 February 2022³

Age Band	12 January – 18 January	19 January – 25 January	26 January – 01 February	02 February – 08 February
Under 18	133	122	119	96
18-29	57	51	37	48
30-39	80	74	56	54
40-49	67	53	63	72
50-54	27	39	30	26
55-59	63	44	29	25
60-64	68	55	27	44
65-69	64	54	46	37
70-74	76	51	47	42
75-79	69	70	47	48
80+	204	172	140	115
Total	908	785	641	607

Source: RAPID (Rapid and Preliminary Inpatient Data)

3. Please refer to [Appendix 3 – Hospital Admissions Notes](#) for explanatory notes regarding RAPID Hospital Admissions.

In the latest week, there has been a 5% decrease in the number of new admissions compared to the previous week, with those aged 80+ years having the highest number of admissions. Also, in the latest week approximately 47% of the hospital admissions related to patients aged 60+.

Table 3 below shows the 'with' COVID-19 hospital admissions by length of stay (LOS), which is another important indicator of measuring the severity of COVID-19.

In the latest week ending 01 February 2022, in all ages, 33% of stays were less than 24 hours, 19% of stays between 24-48 hours and 48% of stays were longer than 48 hours. There was noticeable variation between age groups, with younger age groups having notably shorter stays than the older age groups.

It's important to note that LOS can be influenced by a variety of factors including; age, reason for admission, co-morbidities and hospital pressures.

The analysis presented below in Table 3 has not been adjusted to account for these factors. Additionally, this information is subject to future revisions due to the completeness of discharge information (approximately 8% of records excluded due to missing discharge information).

Table 3: COVID-19 hospital admissions by length of stay, weeks ending 12 January – 01 February 2022⁴


Age Band	12 January - 18 January			19 January - 25 January			26 January - 01 February		
	< 24 Hours	24-48 Hours	>= 48 Hours	< 24 Hours	24-48 Hours	>= 48 Hours	< 24 Hours	24-48 Hours	>= 48 Hours
Under 18	60%	16%	24%	66%	12%	21%	61%	16%	23%
18-29	43%	26%	31%	51%	15%	34%	41%	31%	27%
30-39	42%	22%	36%	47%	23%	30%	42%	27%	31%
40-49	33%	13%	54%	38%	16%	45%	40%	18%	42%
50-54	23%	23%	54%	37%	10%	53%	28%	16%	56%
55-59	20%	16%	65%	13%	13%	75%	22%	17%	61%
60-64	19%	19%	63%	18%	8%	74%	33%	11%	56%
65-69	9%	11%	80%	8%	11%	82%	24%	8%	68%
70-74	17%	4%	80%	22%	9%	69%	6%	22%	72%
75-79	7%	12%	81%	0%	12%	88%	7%	25%	68%
80+	8%	6%	86%	13%	6%	81%	3%	8%	89%
All Ages	29%	15%	56%	35%	13%	52%	33%	19%	48%

4. Length of stay can only be calculated where discharge information has been submitted. Approximately 8% of patients are excluded each week as they are still in hospital or the discharge date has not yet been received. Where a patient has had more than one admission over a 48-hour period, these have been considered as the same day.

In recent months, the proportion of all people who were admitted to hospital within 14 days of a COVID-19 positive test (PCR or LFD) has been declining since early 2021. In the most recent week ending 30 January 2022, there was a proportion of 2% (Figure 7).

The statistics presented in this section reflect the change in case definition (an individual's first positive PCR or LFD from 05 January 22). More information available on the Public Health Scotland website [here](#).

Figure 7: Proportion of weekly cases admitted to hospital within 14 days of a first positive test



Test and Protect

Scotland's approach to contact tracing has continued to adapt throughout the pandemic to reflect changing circumstances, variability in cases, and increasing proportion of the population fully vaccinated since the roll out of the vaccination programme. The most recent [Strategic Framework](#) issued by the Scottish Government in November 2021 sets out how Scotland will continue to adapt now that we are in the phase described as "beyond level zero". That will require a constant review of the associated management information compiled in the weekly report. The information we produce will change over time to reflect the most critical information to help understand, plan and deliver contact tracing at any given point in time.

World Health Organisation (WHO) current guidance on "[Contact tracing in the context of COVID-19](#)" focuses on targeted approaches to contact tracing based on transmission patterns, engaging communities, and prioritising follow-up of high risk cases when it is not possible to identify, monitor and quarantine all contacts. For further information, please refer to [Appendix 2](#).

Please note, PHS has moved to weekly reporting of this data and cumulative data is available in the [interactive dashboard](#). Data for the most recent week, previously included as provisional, is no longer included as this is variable due to cases, which are still open because (either contact tracing is still underway or the NHS Board is still managing the case for a particular reason). Only finalised data will be included within the report going forward.

Index cases

Further background information and definitions are available in [Appendix 4](#).

An **index case** is generated for each positive result with a test date on or after 28 May 2020. This includes tests derived from Scottish and UK Government, as well as self-reported LFD's.

An **individual** is a unique person who has had a positive test. An individual can have multiple positive tests which results in multiple cases within the test and protect system. In these figures, each person is only counted once.

Contact Tracing figures for the week ending 06 February 2022 (based on test date), are detailed in Table 4 below, which provides a recent time trend. A longer time trend is available on the [interactive dashboard](#).

Table 5 provides details of the status of the index cases for each week.

In the week ending 06 February 2022, there were 51,775 Index Cases, of which 38,180 (73.7%) had completed contact tracing by telephone or other digital methods, and a further 6 are in progress (0.01%).

Table 4: Contact Tracing trend information, by week ending

	02 Jan	09 Jan	16 Jan	23 Jan	30 Jan	06 Feb
Total Index Cases ¹	131,762	93,283	54,398	54,687	54,427	51,775
Individuals ²	124,303	87,134	50,767	51,673	51,219	48,445

1. Does not include “Excluded” cases, which are those where a decision has been made that the case should not have been created within the contact tracing system.
2. A count of unique individuals with a positive test. An individual can have multiple positive tests which results in multiple cases within the contact tracing system.

Table 5: Contact Tracing trend information by status, by week ending

Status of cases	02 Jan	09 Jan	16 Jan	23 Jan	30 Jan	06 Feb	Cumulative (from May 2020)
New/ Not yet started ¹	0	0	0	2	0	2	40
% New/ Not yet started	0	0	2	0	2	2	
In progress ²	0	0	0	1	2	6	9
% In progress	0	0	0	0	0	0	
Complete ³	60,869	37,943	30,372	38,544	40,923	38,180	1,007,707
% Complete	46.2	40.7	55.8	70.5	75.2	73.7	
Incomplete ⁴	70,893	55,340	24,024	16,142	13,500	13,587	320,411
% Incomplete	53.8	59.3	44.2	29.5	24.8	26.2	

1. New – New/not yet started cases within the contact tracing system.
2. In progress – The case is still in progress with either the case interview to be completed, or contacts related to the case to be followed up.
3. Complete - The case is complete and all achievable contact tracing has been carried out.
4. Incomplete - Unsuccessful attempts to reach or carry out a case interview via the telephone, or for the index case to provide contacts via digital contact tracing.

Method of Contacting Index Cases

The data within this section are based on the number of **completed cases**, which are recorded in the contact tracing software; these figures are preliminary and may be updated in subsequent publications.

Public Health Scotland works closely with National Services Scotland (NSS) and the Scottish Government to enable local NHS Boards and the National Contact Centre (NCC) to carry out COVID-19 contact tracing effectively. The approach to contact tracing has adapted as restrictions and policy have changed throughout the pandemic in order to best meet the needs of the Scottish population. As numbers of new cases have increased, the method has changed from attempting to phone all new cases and contacts - to prioritising the highest risk cases for telephone calls and sending public health advice by SMS text or email to all others, who have tested positive for COVID-19 and their close contacts.

The introduction of SMS messaging was designed to get public health advice about isolation to cases and contacts as quickly as possible; this is especially pertinent when daily case numbers are very high. The approach was part of a deliberate decision to manage resources through an agreed framework and is in keeping with the evidence-informed advice of the European Centre for Disease Control.

All index cases will receive an initial SMS or email containing Public Health information and advice, which will then be followed by contact either by telephone, additional SMS or email messages containing further Public Health information and advice.

Table 6 below shows a breakdown of the methods used to contact **completed** index cases over time.

Table 6: Contact method used for contact tracing of completed index cases trend information

	02 Jan	09 Jan	16 Jan	23 Jan	30 Jan	06 Feb
Telephone	16,008	19,625	26,263	28,653	28,033	27,159
% Telephone	26.3	51.7	86.5	74.3	68.5	71.1
SMS	44,861	18,318	4,109	9,891	12,890	11,021
% SMS	73.7	48.3	13.5	25.7	31.5	28.9

1. SMS includes those cases deemed low risk and have completed the Co3 online form, every other completed case is categorised as Telephone

In the week ending 06 February 2022, 71.1% of index cases received a telephone call.

Time for a Positive Index Case to be Contact Traced

The data within this section are based on the number of **completed cases**, which are recorded in the contact tracing software; these figures are preliminary and may be updated in subsequent publications.

The three measures shown are:

- the time between a sample being taken and the positive individual being contacted (i.e. interviewed by a contact tracer or completing the online tracing form)
- the time between the record appearing in the CMS and the positive individual being contacted (i.e. interviewed by a contact tracer or completing the online tracing form)
- the time between the record appearing in the CMS and contact tracing being closed (i.e. contacts have been interviewed, attempted to be interviewed or contacted digitally)

These figures are now weekly measures; data are available for previous weeks within the [interactive dashboard](#).

Table 7 and Figure 8 below describes the timeliness of contact tracing by calculating the hours between a test sample being taken and the index case being contacted by Test and Protect either by phone or SMS.

Table 7: Time (hours) between date test sample taken (specimen date) and the positive index case being contacted, for cases completed⁵

Hours taken	Week Ending 06 February 2022		
	Number of Complete Index Cases	% of Total Complete Cases	% of Total Complete & Incomplete Cases
0-24	22,212	58.2	42.9
24-48	10,372	27.2	20.0
48-72	1,749	4.6	3.4
Over 72	495	1.3	1.0
Not recorded* - SMS	533	1.4	1.0
Not recorded* – Phone	2,819	7.4	5.4
Total Complete Cases	38,180	100	
Incomplete Cases	13,587		26.2
Total Complete & Incomplete Cases	51,767		100

⁵ For further information and additional notes on Contact Tracing, please see [Appendix 4 – Contact Tracing](#)

*Improvements into recording of times and dates are being investigated and technical solutions will be identified to reduce the proportion of ‘Not recorded’ cases. This will be implemented February 2022.

Figure 8: Trend in time (hours) between date test sample taken (specimen date) and the positive individual being called for cases completed; by week




Figure 8 shows that more positive cases were contacted over 72 hours after their test sample was taken in June 2021, August 2021 and December 2021, which corresponds with a rise in cases over the same periods.

Table 8: Time (hours) between case created in CMS and the positive individual being contacted^{5, 6}

Hours taken	Week Ending 06 February 2022		
	Number of Complete Index Cases	% of Total Complete Cases	% of Total Complete & Incomplete Cases
0-24	32,589	85.4	63.0
24-48	1,852	4.9	3.6
48-72	213	0.6	0.4
Over 72	183	0.5	0.4
Not recorded* – SMS	533	1.4	1.0
Not recorded* - Phone	2,810	7.4	5.4
Total Complete Cases	38,180	100	
Incomplete Cases	13,587		26.2
Total Complete & Incomplete Cases	51,767		100

5 For further information and additional notes on Contact Tracing, please see [Appendix 4 – Contact Tracing](#)

6 Includes being interviewed by a contact tracer or submitting preliminary information via a CO3 form

*Improvements into recording of times and dates are being investigated and technical solutions will be identified to reduce the proportion of ‘Not recorded’ cases. This will be implemented February 2022.

Table 9: Time (hours) between case created in CMS to its closure^{5, 7}

Hours taken	Week Ending 06 February 2022		
	Number of Complete Index Cases	% of Total Complete Cases	% of Total Complete & Incomplete Cases
0-24	31,975	83.8	61.8
24-48	3,164	8.3	6.1
48-72	532	1.4	1.0
Over 72	341	0.9	0.7
Not recorded* – SMS	384	1.0	0.7
Not recorded* - Phone	1,784	4.7	3.4
Total Complete Cases	38,180	100	
Incomplete Cases	13,587		26.2
Total Complete & Incomplete Cases	51,767		100

5 For further information and additional notes on Contact Tracing, please see [Appendix 4 – Contact Tracing](#)

7 Measured by the time taken to complete the final contact interview for high-risk settings/contacts and those completed via SMS

*Improvements into recording of times and dates are being investigated and technical solutions will be identified to reduce the proportion of ‘Not recorded’ cases. This will be implemented February 2022.

Incomplete index cases

Table 10 and Figure 9 below show the different reasons why an index case is categorised as incomplete (previously referred to as failed) within the contact tracing system.

Incomplete cases are defined as: unsuccessful attempts to carry out a case interview via the telephone, or for the index case to provide contacts via digital contact tracing. This would include scenarios where the mobile/home phone/email address provided by the case was incorrect and no other method of contact could be established; where SMS/telephone call attempts to the case had been made but not been successful in eliciting a response from the index case; where the index case has failed to pass relevant data protection identity checks and where the index case has refused to participate in the contact tracing process.

For operational purposes, some index cases are categorised as incomplete because the telephone process has started, but does not complete for the reasons outlined in Table 10 below. Public Health information is typically sent by SMS to 99% of the incomplete index cases.


Table 10: Number of incomplete index cases by reason

Reason for Incompletion	Week Ending 06 February 2022	
	Number of Index Cases	% of Incomplete Index Cases
Failed ID & verification	131	1.0
No response to call	12,312	90.6
No/incorrect phone number	524	3.9
Refused to provide contact details	32	0.2
Declined to participate / unable to recall contacts	374	2.8
Timed out ¹	214	1.6
Total incomplete cases	13,587	100.0
% incomplete as proportion of all index cases		26.2

1. Timed out cases refers to the closure of cases older than 72 hours. These cases will receive public health information and support signposting via SMS/email communications from Test and Protect. This process allows contact tracing staff to prioritise higher-risk cases and contacts.

In week ending 06 February 2022, 90.6% of incomplete index cases were due to the index case not responding to calls from Test and Protect.

Figure 9: Proportion of reasons for incomplete index cases



Contacts

The Test and Protect system ensures all positive index cases are asked to identify their close contacts, whether they were contacted by telephone and/or SMS. Table 11 below shows the recent trend information of contacts reported to Test and Protect by the index case.

Table 11: Contact Tracing contacts trend information, by week ending

	02 Jan	09 Jan	16 Jan	23 Jan	30 Jan	06 Feb
Total Primary Contacts ¹	106,878	61,756	58,968	72,407	70,934	58,131
Unique Primary Contacts ²	96,630	53,476	49,059	60,029	59,102	50,245
Average number of primary contacts per case	0.8	0.7	1.1	1.3	1.3	1.1

1. Total number of primary contacts recorded in the contact tracing system.

2. Unique number of primary contacts each week. A contact may have been in close contact with multiple index cases.

The average number of primary contacts per case has remained stable over recent weeks.

Contacts not required to self-isolate

Following the Scottish Government announcement on 05 January 2022, from 06 January 2022 fully vaccinated adults and those under the age of 18 years and 4 months do not need to self-isolate as long as they return a negative LFD test result for 7 consecutive days and remain fever free. This applies to both household and non-household contacts. If any of the LFD tests are positive, the contact will be managed as an index case and will not need to book a follow-up PCR to confirm the positive LFD result.

Since the beginning of contact tracing, a small proportion of primary contacts who were successfully contacted were advised they did not need to isolate. Up to 06 February 2022, a total of **3,446** cumulative primary contacts, pertaining to completed index cases, were not advised to self-isolate. This represents **1.1%** of the total **307,234** cumulative primary contacts for which this information is known. Some reasons why contacts do not need to isolate include; children under the age of 16, contact was wearing PPE or did not come into close contact with a positive case.

In the week ending 06 February 2022, of the **50,245** unique contacts recorded, **5,705** (11.4%) went on to test positive (PCR or LFD) within ten days of their contact with an index case.

COVID-19 Vaccine

On 08 December 2020, a COVID-19 vaccine developed by Pfizer BioNTech was first used in the UK as part of national immunisation programmes. The AstraZeneca (Vaxzevria) vaccine was also [approved for use](#) in the national programme, and rollout of this vaccine began on 04 January 2021. Moderna (Spikevax) vaccine was approved for use on 08 January 2021 and rollout of this vaccine began on 07 April 2021. These vaccines have met strict standards of safety, quality and effectiveness set out by the independent Medicines and Healthcare Products Regulatory Agency (MHRA).

For most people, a 2-dose schedule is advised for the vaccines. For the Pfizer BioNTech (Comirnaty) vaccine, the second vaccine dose can be offered between 3 to 12 weeks after the first dose. For the AstraZeneca (Vaxzevria) and Moderna (Spikevax) vaccine, the second dose can be offered 4 to 12 weeks after the first dose.

Information on uptake across the vaccine programme is available on a daily basis via the PHS [COVID-19 Daily Dashboard](#), 5 days a week at 2pm (Monday to Friday). This provides a cumulative picture of the position nationally and locally.

The dashboard provides total uptake nationally with breakdowns by [Joint Committee on Vaccination and Immunisation \(JCVI\)](#) age based cohorts and non-age based cohorts for priority groups 1-9.

The vaccination content of this weekly publication is kept under continual review and specific editions have contained more in-depth analyses of uptake by particular groups or characteristics, including uptake by ethnicity and deprivation category, for teachers, for prisoners and for pregnant women.

COVID-19 Vaccination Uptake


As of 14 February 2022, there has been over 11.9 million Covid-19 vaccine doses administered in Scotland, since the programme began on 08 December 2020.

- 4.43 million people protected through their first dose of the COVID-19 vaccination; 93.6% of those aged 18 and over and 92.2% of those aged 12 and over.
- 4.14 million people provided with further protection by receiving their second dose, of these, 89.8% are aged 18 and over and 86.6% of those aged 12 and over.
- 3.35 million people have received their booster/dose 3, of these, 75.2% are aged over 18 and over and 70.1% of those aged 12 and over.

More detailed age information can be in Figure 10

Daily Vaccination uptake information is available via the [PHS Covid Daily Dashboard](#).

Figure 10: Covid-19 Vaccine uptake – percentage coverage by age group in Scotland



COVID-19 Vaccine Wastage

Given the scale of the Covid-19 vaccination programme, some vaccine wastage has been unavoidable for a variety of reasons including logistical issues, storage failure and specific clinical situations.

The initial planning assumption for the vaccination programme was that there would be around 5% vaccine wastage. Table 12 below shows the trend of the percentage of vaccines wasted by calendar month between August 21 – January 2022.

In January 2021, the percentage of vaccines wasted was 2.2%. The top reasons for doses wasted in this month were: excess stock (55%), expired shelf life of stock (31%) other reasons (14%).

Excess stock is defined: Where a vaccination team reach the end of an allotted shift or job, and have surplus vaccines that cannot be returned to stock, or used before it expires. This includes any unused doses in opened vials at the end of a clinic.

Table 12: Number of COVID-19 Vaccination doses wasted by Month^{1,2,3,4,5}

Measure	Aug-21	Sept-21	Oct-21	Nov-21	Dec-21	Jan-22
Number of doses administered ¹	568,203	303,931	884,381	1,017,131	1,369,363	471,925
Number of doses wasted ²	17,457	10,089	9,116	8,605	8,608	10,783
Percentage wasted ³ (%)	2.9	3.2	1.0	0.8	0.6	2.2

Source: NSS Service Now, COVID-19 Vaccine Wastage.

Data correct at 15/2/22

1. The number of vaccine doses administered (all doses).
2. The total number of vaccine doses which could not be administered and therefore wasted.
3. % Wasted is measured as:
(Number of Doses Wasted x 100)/(Number of Doses Wasted + Administered)
4. Excludes GP practice information.
5. Excludes wastage from clinical trials

COVID-19 Vaccine Certification

To show COVID-19 vaccine status, there are a number of options and individuals can choose to use one or more of these:

- Use the NHS Covid Status App
- Request a paper copy of your COVID-19 Status
- Download a PDF copy of your COVID-19 Status

The NHS Covid Status App was launched on 30 September 2021. It is free and offers digital proof of vaccination via a QR code for each vaccination received. You can request a vaccine certificate if you are aged 12 and over and have been vaccinated in Scotland. The record will not show any vaccinations given outside of Scotland.

- As of midnight 12 February 2022, the NHS Covid Status App has been downloaded 2,643,193 times. It is important to note a single user may choose to download the App on multiple devices, so this figure does not represent unique individuals
- Between 03 September 2021 (introduction of QR codes) and midnight 12 February 2022
 - 828,471 paper copies of COVID-19 Status have been requested. This may not represent unique users if an individual requests a second copy (for example if they have lost their paper copy)
 - 1,947,161* PDF versions of COVID-19 Status have been downloaded. This provides a measure of the total number of times a new QR code has been generated via PDF. An individual can generate more than one successful QR code so the figure does not represent unique users

*01, 02 and 03 October data for PDFs is missing due to a technical error, we can reasonably estimate that there were 35,000 – 45,000 PDFs successfully generated PDFs in total for those three days.

COVID-19 Cases, Acute Hospitalisations, and Deaths by Vaccine Status

Vaccine Surveillance

Public Health Scotland has a [COVID-19 vaccine surveillance strategy](#) to monitor the effectiveness, safety and impact of all approved COVID-19 vaccines in Scotland. The key measure of the success of the vaccination programme in preventing infection, hospitalisations and deaths is vaccine effectiveness. The summary data presented in this section record the total number of COVID-19 cases, COVID-19 related acute hospital admissions and confirmed COVID-19 deaths by their vaccination status.

Please note that this section only includes PCR confirmed COVID-19 cases, COVID-19 related acute hospital admissions and confirmed COVID-19 related deaths. Lateral flow device (LFD) cases are not included in this section and therefore will result in an underestimation of recent case rate trends.

PLEASE READ BEFORE REVIEWING THE FOLLOWING TABLES AND FIGURES

Please note that due to a coding error, the previous four published weekly reports (from 19 January 2022 to 09 February 2022) included PCR-confirmed cases for ages 15 years old and over for the case numbers and age-standardised case rates, rather than 10 years old and over as reported. Data in this report as been updated to include 10 years and over and previous reports will also be updated to reflect the correction.

NOTICE OF CHANGE

From 16 February 2022, Public Health Scotland (PHS) will no longer report COVID-19 cases, hospitalisations, and deaths by vaccination status on a weekly basis. PHS will continue to provide updates from the latest scientific analyses and reports on the effectiveness of COVID-19 vaccines. An explanation behind this decision can be found below.

Since July 2021, Public Health Scotland (PHS) have reported COVID-19 cases, hospitalisations, and deaths by vaccination status in the weekly COVID-19 statistical report. These data were first published to help monitor the impact of the vaccination programme on the pandemic, impact on the NHS, and to help understand where to prioritise vaccination delivery. While PHS has stated that the data in the report should not be used as a measure of vaccine effectiveness, PHS is aware of inappropriate use and misinterpretation of the data when taken in isolation without fully understanding the limitations described below.

Due to the increasing risk of misinterpretation from growing complexities as the COVID-19 pandemic enters its second year (as described below), PHS has taken the decision to no longer report COVID-19 cases, hospitalisations and deaths by vaccination status on a weekly basis. PHS is currently reviewing the content and frequency of reporting this information. PHS will continue to communicate up to date and high quality research on COVID-19 vaccines. [Evidence suggests](#) the COVID-19 vaccine is effective at reducing the risk of a severe outcome, such as hospitalisation when infected, however no vaccine is 100% effective and some COVID-19 hospitalisations and deaths may still occur, particularly in [vulnerable populations](#). PHS, in collaboration with partners such as the [EAVE II](#)

[consortium](#), will continue supporting comprehensive scientific studies evaluating the risks and reasons for severe COVID-19 in vaccinated populations and will communicate its findings as and when appropriate.

NOTICE OF CHANGE: Reasons why it is inappropriate to continue publishing case rates

Recent changes in testing behaviour and policies, including the fact that [asymptomatic people no longer need a COVID-19 PCR test](#), limits our ability to robustly identify and monitor COVID-19 cases by vaccination status. **This means we cannot confidently compare recent case trends to historical data.**

COVID-19 rates do not account for potential differences between populations such as behaviour towards social distancing, underlying health issues, or approaches to testing and how much they contribute to COVID-19 rates. For example, [a study](#) found that people with two doses of vaccine were more likely to test themselves for COVID-19 compared to those unvaccinated or with one dose of COVID-19 vaccine. This means that unvaccinated people may be less likely to test and report as a case resulting in lower infection rates among the unvaccinated. **Comparison of case rates is therefore now inappropriate.**

The population of Scotland is relatively small compared to other countries such as England and the United States, and with a high vaccination rate, systematic underlying differences between the unvaccinated, partially vaccinated and booster populations become more evident sooner than bigger countries that may have a larger unvaccinated population. In Scotland most people are vaccinated and have received a booster, this results in a small number of people who are unvaccinated or have had one dose or two doses. Small increases in case numbers in a small population have a bigger impact on case rates than small increases in a larger population (e.g. boosted) where such differences are less noticeable due to the sheer number of people. **Comparison of case rates is therefore inappropriate.**

Vaccinations have not been given to the entire population at the same time with [certain groups given priority before others](#). People are on different timelines since vaccination, dose schedules (e.g. some groups are not eligible for a booster) and vaccine types creating difficulties when making broad comparisons across the population by vaccine status. A number of people have not completed or exceeded their recommended dose schedule. This leads to [reduced protection](#) from the vaccine over time. Such differences between people impacts how we interpret the rates by vaccine status, for example the level of vaccine protection is expected to be higher in someone who had their first dose four weeks ago compared to another who had their first dose one year ago and didn't have the required second dose as per the vaccination schedule. **Comparison of case rates is therefore inappropriate.**

Individuals who have not completed their vaccine schedule [may be more susceptible](#) to a severe outcome and could result in higher COVID-19 case, hospitalisation and death rates in the first and second dose vaccine groups. For example, some of the older individuals who have exceeded the recommended time will have not received their next vaccine dose because of frailty or ill health. They may, therefore, be more likely to be hospitalised or die if they get COVID-19. **Comparison of hospitalisation and death rates is therefore inappropriate.**

There is a lot uncertainty about the number of people in the unvaccinated population because we do not know if individuals are still resident in Scotland. This is unlike the vaccinated population who have had recent contact with the NHS due to vaccination. As case rates require accurate population estimates, the uncertainty in the unvaccinated population will result in an underestimate of case rates. **Comparison of case rates is therefore inappropriate.** This is explained in more detail in [our blog](#).

NOTICE OF CHANGE: Summary

Public Health Scotland is committed to providing high quality public health science, communication, and transparency. Vaccine effectiveness studies are needed to look at this complicated data and evaluate how well the COVID-19 vaccines are working. For the reasons outlined above, we will no longer provide a weekly summary of COVID-19 cases, hospitalisation, and deaths by vaccine status, but focus our attention to vaccine effectiveness reporting.

PLEASE READ BEFORE REVIEWING THE FOLLOWING TABLES AND FIGURES

Interpretation of data

There is a large risk of misinterpretation of the data presented in this section due to the complexities of vaccination data. [A blog post](#) by the UK Health Security Agency (UKHSA), formerly Public Health England (PHE), provides a comprehensive explanation of the biases and potential areas for misinterpretation of such data. **They state that a simple comparison of COVID-19 case rates in those who are vaccinated and unvaccinated should not be used to assess how effective a vaccine is in preventing serious health outcomes, because there are a number of differences between the groups, other than the vaccine itself, and these biases mean that you cannot use the rates to determine how well the vaccines work.**

Below are examples of some of the complexities and biases that need to be taken into consideration when interpreting these data. This is explained in more detail in [our blog](#) on the PHS website.

Vaccinated individuals can still be infected with COVID-19

In Scotland, there has been a very high uptake of the COVID-19 vaccine. As of 04 February 2022, 89.7% of 18 years old and over have received a second dose and 74.4% have received a third dose or booster of COVID-19 vaccine. No vaccine is 100% effective and it's expected that cases, hospitalisations, and deaths from COVID-19 will occur in the vaccinated population as well as the unvaccinated population.

The [current evidence](#) suggests that you may test positive for COVID-19, or be reinfected even if you are vaccinated, especially since the emergence of the Omicron variant in the UK. The major benefit of vaccination against Omicron is to protect from severe disease. [Follow public health guidance](#) and test yourself if you develop any COVID-19 like symptoms.

COVID-19 vaccines protect most people against severe outcomes, but some people will still get sick because no vaccine is 100% effective

[Evidence suggests](#) the COVID-19 vaccines are over 75% effective at preventing a severe outcome of COVID-19. COVID-19 [hospitalisations](#) and [deaths](#) are strongly driven by older age, with most deaths occurring in those over 70 years old and having multiple other illnesses. But overall, you are less likely [to be hospitalised if you are vaccinated with a booster](#).

Data and rates presented in this section are not a measure of vaccine effectiveness

Vaccine effectiveness is a scientific method used to measure how well a vaccine protects people against outcomes such as infection, symptoms, hospitalisation and death in the 'real-world'. Unlike case rates, vaccine effectiveness analysis accounts for potential biases in the data and risk factors such as age, sex, prior infection, co-morbidities, socio-economic status and time since vaccination. This method is the most robust way to measure if a vaccine is working.

The data and rates presented in this section do not account for biases and risk factors and should not be used to measure vaccine effectiveness. We include links to vaccine effectiveness studies below.

There are likely to be systematic differences and biases between the vaccinated and unvaccinated groups, such as behaviour, vulnerability and previous infection, that are unaccounted for when comparing rates. As most of the population is vaccinated, these differences become more evident and could create bias in case/hospitalisation/death rates between vaccinated and unvaccinated population. For example, **people who are vaccinated may be more likely to follow government guidance such as regular testing and reporting for COVID-19.** [A study](#) found that **people with two doses of vaccine had higher intentions to get tested for COVID-19 compared to those unvaccinated or having had one dose of COVID-19 vaccine.** These differences in behaviour for vaccinated individuals makes them more likely to be identified as a case than unvaccinated people, resulting in higher case rates in the vaccinated population.

Vaccine effectiveness wanes over time

COVID-19 vaccine effectiveness [waned over time](#). Within the first and second dose population there will be a number of individuals that will have exceeded the recommended time for their next vaccine dose. These people [may be more susceptible](#) to a severe outcome and could result in higher COVID-19 case, hospitalisation and death rates in the first and second dose vaccine groups. For example, some of the older individuals who have exceeded the recommended time will have not received their next vaccine dose because of frailty or ill health. They are, therefore, more likely to be hospitalised or die if they get COVID-19.

Difficulty knowing who is in the unvaccinated population

There is a lot of uncertainty about who is in the unvaccinated population, which makes interpretation of COVID-19 rates in this group difficult. Everyone in Scotland who is registered with a GP is assigned a unique Community Health Index (CHI) number. This number is how we estimate Scottish residents for the analysis in this section, however, it has its limitations when people leave Scotland and do not inform their GP, resulting in an overestimate of Scottish residents. To try and account for this issue, we also use COVID-19 vaccination records to calculate the number of people in the vaccinated population. This helps determine whether people are still resident in Scotland, however, those in the unvaccinated population have not had recent contact with a vaccination centre and therefore we have to rely on GP records, which are likely to be more out of date than vaccine records. This is explained in more detail in [our blog](#).

Vaccine effectiveness summary

[UKHSA publish](#) a summary of current vaccine effectiveness in their weekly COVID-19 vaccine surveillance report. [The latest evidence](#) can be found below, which suggests the COVID-19 booster/third dose vaccine lowers your risk of a severe outcome compared to the second dose.

The Office for National Statistics has also published research into the [risk of testing positive](#) for COVID-19 by vaccination status, [impact of Delta](#) on viral burden and vaccine effectiveness, and the [risk of death](#) by vaccination status.

Figure 11: [Summary of evidence](#) on vaccine effectiveness against different outcomes for Omicron (all vaccines combined)

	Dose 2			Dose 3		
	0-3 months	4-6 months	6+ months	0-3 months	4-6 months	6+ months
Infection	Insufficient data	Insufficient data	Insufficient data	Insufficient data	Insufficient data	Insufficient data
Symptomatic disease	25-70%	5-30%	0-10%	50-75%	40-50%	Insufficient data
Hospitalisation	65-85%	55-65%	30-35%	80-95%	75-85%	Insufficient data
Mortality	Insufficient data	Insufficient data	40-70%	85-99%	Insufficient data	Insufficient data

Figure 12: [Summary of evidence](#) on vaccine effectiveness against different outcomes for Delta (all vaccines combined)

	Dose 2			Dose 3		
	0-3 months	4-6 months	6+ months	0-3 months	4-6 months	6+ months
Infection	65-80%	50-65%	Insufficient data	Insufficient data	Insufficient data	Insufficient data
Symptomatic disease	65-90%	45-65%	40-60%	90-99%	90-95%	Insufficient data
Hospitalisation	95-99%	80-90%	70-85%	95-99%	Insufficient data	Insufficient data
Mortality	95-99%	90-95%	80-99%	95-99%	Insufficient data	Insufficient data

Figure 13: Confidence rating scale

High Confidence	Evidence from multiple studies which is consistent and comprehensive
Medium Confidence	Evidence is emerging from a limited number of studies or with a moderately level of uncertainty
Low Confidence	Little evidence is available at present and results are inconclusive

Summary of key results

- Case rates have declined and subsequently plateaued in the last four weeks from 15 January 2022 to 11 February 2022. Caution should be taken when interpreting recent case trends due to the change in [policy](#) from 06 January 2022 where an asymptomatic individual who tests positive via a LFD test is not required to take a confirmatory PCR test. This section of the report only includes PCR confirmed COVID-19 cases and does not include cases confirmed by a LFD test.
- In the most recent four weeks, from 15 January 2022 to 11 February 2022, the age-standardised rate of hospital admissions per 100,000 were 3.0 to 3.7 times lower in individuals with their third dose or booster dose of vaccine compared to unvaccinated individuals or have received one or two doses of a COVID-19 vaccine.
- In the four weeks from 08 January 2022 to 04 February 2022, in an age-standardised population, the death rate in individuals that received a booster or 3rd dose of a COVID-19 vaccine was between 4.6 and 9.5 times lower than individuals who are unvaccinated or have only received one or two doses of a COVID-19 vaccine.

Overall results of COVID-19 cases and hospitalisations, and deaths by vaccination status

COVID-19 cases by vaccination status

[Analyses from Scotland](#) show that the booster and third dose of the COVID-19 vaccines are associated with 57% reduced risk of symptomatic infection with the Omicron variant compared to those who are more than 25 weeks post-second dose of COVID-19 vaccine.

[Analysis](#) of vaccine effectiveness against symptomatic disease with the Omicron variant suggests a lower impact than the Delta variant. Although lower, this is still a substantial vaccine effect, and effectiveness is higher after a booster than after one or two doses of vaccine. For the latest vaccine effectiveness estimates from UKHSA, please see the section [Vaccine effectiveness summary](#) above.

Table 13: PCR-confirmed COVID-19 age-standardised case rate per 100,000 individuals by vaccine status, seven-day rolling average from 15 January 2022 to 11 February 2022

	Unvaccinated			1 Dose*		
Week	No. tested positive by PCR	Population	Age-standardised case rate per 100,000 with 95% confidence intervals	No. tested positive by PCR	Population	Age-standardised case rate per 100,000 with 95% confidence intervals
15 January - 21 January 2022	5,320	976,941	439.48 (416.28 - 462.67)	1,962	318,871	481.31 (443.73 - 518.89)
22 January - 28 January 2022	4,956	970,309	381.51 (362.44 - 400.59)	1,664	302,843	422.99 (388.50 - 457.49)
29 January - 04 February 2022	4,757	962,727	393.55 (374.97 - 412.13)	1,444	275,689	383.99 (353.98 - 413.99)
05 February - 11 February 2022	3,834	956,449	340.79 (321.48 - 360.10)	1,152	262,647	343.90 (315.71 - 372.08)
	2 Doses*			Booster or 3 Doses		
Week	No. tested positive by PCR	Population	Age-standardised case rate per 100,000 with 95% confidence intervals	No. tested positive by PCR	Population	Age-standardised case rate per 100,000 with 95% confidence intervals
15 January - 21 January 2022	6,522	934,811	617.62 (596.83 - 638.40)	10,772	3,070,303	428.00 (401.48 - 454.51)
22 January - 28 January 2022	5,411	855,982	569.85 (548.82 - 590.87)	11,123	3,170,692	446.83 (415.17 - 478.49)
29 January - 04 February 2022	5,079	830,753	525.86 (505.59 - 546.13)	12,052	3,229,937	500.21 (474.93 - 525.49)
05 February - 11 February 2022	5,201	809,783	549.69 (529.08 - 570.31)	13,833	3,270,226	527.98 (508.76 - 547.21)

* 1 Dose and 2 Dose populations include individuals who have exceeded the recommended dose schedule and may be subject to vaccine waning. Data in this table should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.

Vaccination status is determined as at the date of PCR specimen date according to the definitions described in Appendix 6. The data displayed within the greyed-out section are considered preliminary and are subject to change as more data is updated. Age-standardised case rates are per 100,000 people per week, standardised to the 2013 European Standard Population (see Appendix 6).

Please note that due to a coding error, the above table for the previous four published weekly reports (from 19 January 2022 to 09 February 2022) included PCR-confirmed cases for ages 15 years old and over for the case numbers and age-standardised case rates, rather than 10 years old and over as reported. Table 13 above contains the corrected numbers and previous reports will also be corrected.

To account for the different age distribution of individuals in each vaccine status, age standardised rates are used in Table 13. These are adjusted to only include individuals 10 years old and over and are calculated by combining rates from different age groups relative to the European standard age distribution population. The calculations have associated 95% confidence intervals shown in the shaded areas of the figure. Smaller populations have wider associated confidence intervals whereas larger populations have narrower associated confidence intervals.

The number of PCR confirmed COVID-19 cases and case rates have plateaued in the last week from 05 February 2022 to 11 February 2022. **Caution should be taken when interpreting recent case trends due to the change in policy from 06 January 2022 where an asymptomatic individual who tests positive via a LFD test is not required to take a confirmatory PCR test.** Individuals may test positive for COVID-19 even if vaccinated. However, the data in Table 13 does not account for severity of the case such as presence of symptoms and may include a number of asymptomatic individuals. Current evidence suggests that the vaccine is very effective at preventing hospitalisations and deaths. The rates in Table 13 should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors. For more information, please see the [Interpretation of data](#) section above.


COVID-19 related acute hospital admissions by vaccine status

[A number of studies](#) have estimated vaccine effectiveness against hospitalisation and have found high levels of protection against hospitalisation with all vaccines against the Alpha variant. [A paper](#) observed effectiveness against hospitalisation of over 90% with the Delta variant with all three COVID-19 vaccines including AstraZeneca (Vaxzevria), Pfizer-BioNTech (Comirnaty), and Moderna (Spikevax). In most groups there was relatively limited waning of protection against hospitalisation over a period of at least five months after the second dose.

Vaccine effectiveness against hospitalisation with the Omicron variant is slightly lower than the Delta variant, but the booster or third dose of COVID-19 vaccine still provides excellent protection against COVID-19 related hospitalisation. For the latest vaccine effectiveness estimates from UKHSA please see the section [Vaccine effectiveness summary](#) above.

From 10 May 2021 to 11 February 2022, there were a total of 1,458,140 acute hospital admissions for any cause, of which 34,523 were associated with a COVID-19 PCR positive test 14 days prior, on admission, the day after admission or during their stay. Using the 90-day exclusion criteria between positive COVID-19 PCR tests associated with an acute hospital admission, 38,700 individuals were admitted to hospital, of which 117 were readmitted more than 90 days after their first admission.

Figure 14: Seven-day rolling average on a log₁₀ scale: acute hospital admissions where the individual had a COVID-19 positive PCR test 14 days prior, on admission or during their stay in hospital, compared to all acute hospital admissions, 10 May 2021 to 11 February 2022



Data in this figure should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.

Data displayed are on a log₁₀ scale. The data displayed within the greyed-out section are considered preliminary and are subject to change as more data is updated.

COVID-19 related acute hospital admissions have increased to similar levels observed in September 2021 and are now declining again. COVID-19 related acute hospital admissions are small relative to all acute hospitalisations.

Table 14: Age-standardised rate of acute hospital admissions where an individual had a COVID-19 positive PCR test up to 14 days prior, on admission, or during their stay in hospital, by week and vaccination status, 15 January 2022 to 11 February 2022

	Unvaccinated			1 Dose*		
Week	No. hospitalised	Population	Age-standardised hospitalisation rate per 100,000 (95% confidence intervals)	No. hospitalised	Population	Age-standardised hospitalisation rate per 100,000 (95% confidence intervals)
15 January - 21 January 2022	114	794,442	31.50 (20.99 - 42.00)	45	230,407	39.87 (22.86 - 56.88)
22 January - 28 January 2022	99	790,408	30.16 (18.90 - 41.43)	30	221,664	29.04 (14.32 - 43.76)
29 January - 04 February 2022	93	785,164	32.15 (20.64 - 43.65)	25	211,760	17.88 (8.04 - 27.71)
05 February - 11 February 2022	64	780,982	15.10 (7.88 - 22.31)	15	207,261	10.90 (3.05 - 18.76)
	2 Doses*			Booster or 3 Doses		
Week	No. hospitalised	Population	Age-standardised hospitalisation rate per 100,000 (95% confidence intervals)	No. hospitalised	Population	Age-standardised hospitalisation rate per 100,000 (95% confidence intervals)
15 January - 21 January 2022	151	917,126	37.89 (29.14 - 46.63)	375	3,069,278	10.75 (9.51 - 11.99)
22 January - 28 January 2022	116	829,249	34.19 (25.48 - 42.91)	317	3,169,578	8.44 (7.47 - 9.41)
29 January - 04 February 2022	93	785,327	25.10 (17.41 - 32.80)	281	3,228,702	7.47 (6.54 - 8.40)
05 February - 11 February 2022	89	754,551	27.30 (18.81 - 35.79)	263	3,268,885	7.22 (6.29 - 8.15)

* 1 Dose and 2 Dose populations include individuals who have exceeded the recommended dose schedule and may be subject to vaccine waning. Data in this table should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.

Vaccination status is determined as at the date of positive PCR test according to the definitions described in Appendix 6. The data displayed within the greyed-out section are considered preliminary and are subject to change as more data is updated. Age-standardised hospitalisation rates are per 100,000 people per week, standardised to the 2013 European Standard Population (see Appendix 6).

To account for the different age distribution of individuals in each vaccine status, age-standardised hospitalisation rates are reported in Table 14 and Figure 15. To align with the 2013 European Standard Population, 15 years olds are now included in the number of acute hospital admissions and the population denominator. This means that data presented in Table 14 may differ from previous weeks reports which only included hospitalised individuals and denominators for 16 years and older. The rates in Table 14 should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors. For more information, please see the [Interpretation of data](#) section above.


In the past four weeks, from 15 January 2022 to 11 February 2022, the age-standardised rate of hospital admissions per 100,000 were 3.0 to 3.7 times lower in individuals with their third dose or booster dose of vaccine compared to unvaccinated individuals or have received one or two doses of a COVID-19 vaccine.

It is important to highlight the wide 95% confidence intervals for the unvaccinated, one dose and two dose age-standardised COVID-19 related acute hospital admission rates. Wide confidence intervals are the result of small populations in these groups as the majority of Scotland has received a COVID-19 booster vaccine. These rates should be interpreted with caution.

From 15 January 2022 to 11 February 2022, of the post second dose acute COVID-9 related hospital admissions in Table 14, approximately 73% received their second dose of vaccine more than 6 months prior to a COVID-19 PCR positive test.

Please note that these statistics do not differentiate between individuals in hospital with COVID-19 illness requiring hospitalisation compared to those in hospital for other reasons (e.g. routine operations) for whom COVID-19 was identified incidentally through testing but they are not requiring hospitalisation because of their COVID-19 symptoms.

Figure 15: Age-standardised hospitalisation rate of acute hospital admissions where an individual had a COVID-19 positive PCR test up to 14 days prior, on admission, or during their stay in hospital, per 100,000 individuals by vaccination status, seven-day rolling average from 10 May 2021 to 11 February 2022



1 Dose and 2 Dose populations include individuals who have exceeded the recommended dose schedule and may be subject to vaccine waning. Data in this figure should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.


Vaccination status is determined as at the date of positive PCR test according to the definitions described in Appendix 6. The data displayed within the greyed-out section are considered preliminary and are subject to change as more data is updated. 95% confidence intervals are shown as the shaded regions. Age-standardised hospitalisation rates are per 100,000 people per week, standardised to the 2013 European Standard Population (see Appendix 6).

To account for the different age distribution of individuals in each vaccine status, age-standardised hospitalisation rates are reported in Table 14 and Figure 15. To align with the 2013 European Standard Population, 15 years olds are now included in the number of acute hospital admissions and the population denominator. This means that data presented in Table 14 may differ from previous weeks reports which only included 16 years and older. The rates in Figure 15 should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors. For more information, please see the [Interpretation of data](#) section above.

In the past four weeks, from 15 January 2022 to 11 February 2022, the age standardised rate of acute hospital admissions for individuals with a booster or third dose remains lower compared to people unvaccinated or who have only received one or two doses of a COVID-19 vaccine.

Note that the peak in the booster or third dose category around 11 October 2021 is likely due to a small number of individuals vaccinated and the prioritisation of the booster/third dose to the clinically extremely vulnerable at the beginning of the booster programme.

Figure 16: Seven-day rolling average COVID-19 related acute hospital admissions by vaccination status and by age group, 10 May 2021 to 11 February 2022



Data in this figure should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.

Vaccination status is determined as at the date of positive PCR test according to the definitions described in Appendix 6. Patient age is determined as their age the date of admission. The data displayed within the greyed-out section are considered preliminary and are subject to change as more data is updated.

Overall, the highest rates of acute COVID-19 related hospital admissions were in the oldest age groups. In groups where a very large proportion of individuals have been vaccinated (such as individuals over age 70), any small changes in COVID-19 related acute hospital admissions will result in a larger change shown in the graph, for example in the over 70 dose 1 vaccinated group. These changes tend to be more ‘step like’ and less smooth.

Confirmed COVID-19 deaths by vaccination status

COVID-19 vaccines are estimated to significantly reduce the risk of mortality for COVID-19, however a small number of COVID-19 deaths are still expected in vaccinated people, especially in vulnerable individuals where the vaccine or the immune response may not have been effective. Evidence has shown that vaccination is highly effective in protecting against death from coronavirus (COVID-19).

[A recent paper](#) examined the effectiveness of the Pfizer (Comirnaty) booster vaccine and found that adults who had received their booster dose five months after their second dose had 90% lower risk of mortality due to COVID-19 than adults who hadn't received their booster dose five months after their second dose.

For the latest vaccine effectiveness estimates from UKHSA, please see the section [Vaccine effectiveness summary](#) above.

Table 15: Number of confirmed COVID-19 related deaths by vaccination status at time of test and age-standardised mortality rate per 100,000, 08 January 2022 to 04 February 2022

	Unvaccinated			1 Dose*		
Week	No. of deaths	Population	Age Standardised Mortality Rate per 100,000 with 95% confidence intervals	No. of deaths	Population	Age Standardised Mortality Rate per 100,000 with 95% confidence intervals
08 January - 14 January 2022	17	1,542,795	11.94 (4.40 - 19.48)	3	339,017	4.79 (0.00 - 11.14)
15 January - 21 January 2022	18	1,538,621	7.46 (2.70 - 12.22)	6	318,899	12.12 (1.58 - 22.67)
22 January - 28 January 2022	13	1,531,988	10.68 (3.82 - 17.54)	2	302,872	2.48 (0.00 - 5.96)
29 January - 04 February 2022	13	1,524,406	10.95 (3.40 - 18.50)	4	275,718	8.57 (0.00 - 17.35)
	2 Doses*			Booster or 3 Doses		
Week	No. of deaths	Population	Age Standardised Mortality Rate per 100,000 with 95% confidence intervals	No. of deaths	Population	Age Standardised Mortality Rate per 100,000 with 95% confidence intervals
08 January - 14 January 2022	34	999,502	14.37 (9.15 - 19.59)	72	2,982,479	1.52 (1.16 - 1.87)
15 January - 21 January 2022	31	934,811	16.15 (10.27 - 22.03)	77	3,070,304	1.60 (1.24 - 1.96)
22 January - 28 January 2022	26	855,982	14.87 (8.82 - 20.93)	70	3,170,693	1.43 (1.09 - 1.77)
29 January - 04 February 2022	17	830,753	11.68 (6.05 - 17.32)	73	3,229,938	1.50 (1.15 - 1.85)

* 1 Dose and 2 Dose populations include individuals who have exceeded the recommended dose schedule and may be subject to vaccine waning. Data in this table should not be used as a measure of vaccine effectiveness due to unaccounted for biases and risk factors in different populations. For more information, please see the [Interpretation of data](#) and [Vaccine effectiveness summary](#) sections above.

Vaccination status is determined as at the date of positive PCR test according to the definitions described in Appendix 6. A confirmed COVID-19 related death is defined as an individual who has tested positive by PCR for SARS-CoV-2 at any time point and has COVID-19 listed as an underlying or contributory cause of death on the death certificate. Age-standardised mortality rates per 100,000 people per week, standardised to the 2013 European Standard Population (see Appendix 6). This definition is for the purposes of evaluating the impact of the COVID-19 vaccine on confirmed COVID-19 deaths. The numbers reported in this section may differ from other published COVID-19 death data. Data are based on date of registration. In Scotland deaths must be registered within 8 days although in practice, the average time between death and registration is around 3 days. More information on days between occurrence and registration can be found on the NRS website.

To account for differences in population size and age of the vaccination status groups over time, age-standardised mortality rates were calculated for deaths where COVID-19 was listed as an underlying or contributory cause of death on the death certificate (Table 15). In the four weeks from 08 January 2022 to 04 February 2022, in an age-standardised population, the death rate in individuals that received a booster or 3rd dose of a COVID-19 vaccine was between 4.6 and 9.5 times lower than individuals who are unvaccinated or have only received one or two doses of a COVID-19 vaccine.

It is important to highlight the wide 95% confidence intervals for the unvaccinated, one dose and two dose age-standardised COVID-19 related acute hospital admission rates. Wide confidence intervals are the result of small populations in these groups as the majority of Scotland has received a COVID-19 booster vaccine. These rates should be interpreted with caution.

From 29 December 2020 (21 days after the start of the vaccination programme in Scotland to account for protection to develop after the first dose) to 04 February 2022, there have been 6,268 confirmed COVID-19 related deaths with a positive PCR result and where COVID-19 was recorded as an underlying or contributory cause on the death certificate.

In Scotland, from the beginning of the COVID-19 vaccination programme over 3.3 million individuals have received a booster or third dose of a COVID-19 vaccine. In the week from 29 January to 4 February 2022, 74 individuals tested positive by PCR for SARS-CoV-2 more than fourteen days after receiving their booster or third dose of COVID-19 vaccine and subsequently died with COVID-19 recorded as underlying or contributory cause of death. The majority of these individuals had several comorbidities, and the mean age was 82 years old (IQR 75 to 91 years old).

In the week from 29 January to 4 February 2022, the majority of the 17 post second dose confirmed COVID-19 related deaths Table 15 received their second dose of COVID-19 vaccine more than six months prior to their death. The majority of these individuals had several comorbidities, and the mean age was 83 years old (IQR 79 to 91 years old).

Hospital/ Wider System Pressures

NHS services across NHS Scotland are subject to increased demand during the winter period. The information presented in this section aims to support the reader in drawing insights from a wider range of existing metrics around COVID-19 and winter pressures.

Unscheduled Care

As individuals in Scotland make contact with Unscheduled Care Services, data about who they are, where they have come from, what is wrong with them and what happens to them are collected, mainly to inform their care. This provides a good picture of the potential unscheduled care journeys that an individual may travel through.

Pressures on unscheduled care services are a familiar sight during the winter. Increased incidence of respiratory infections, alongside an increased acuity of illness and demands on primary care leads to increased demand on unscheduled care.

NHS inform is Scotland's digital health and care resource, providing the up to date standardised information on COVID-19 from a health perspective. Information is provided in a range of languages and alternative formats (www.nhsinform.scot/coronavirus).

Additional information can be found on the [wider impacts dashboard](#) and also in our [interactive dashboard](#).

NHS 24

During COVID-19, there has been a rapid reconfiguration of primary and community care services. As part of this NHS 24's 111 service has been reconfigured as an in-hours (as well as out-of-hours) route for COVID-19 triage for rapid access to care via local COVID-19 assessment hubs. In addition to this, from 1st December 2020, the national Redesign of Urgent Care Programme introduced new pathways from NHS 24 to Flow Navigation Centres, with the aim of reducing the numbers of people attending A&E and diverting to more appropriate care closer to home. This is available as part of a 24/7 service, further increasing NHS 24 in-hours activity (Monday to Friday, 8am to 6pm).

Information on COVID-19 related contacts to NHS24 and the Coronavirus Helpline are presented in our [interactive dashboard](#), which supplements this report.

Primary Care Out of Hours (OOH)

Across Scotland, NHS Boards provide Primary Care Out of Hours (OOH) services for patients' when their registered GP practice is closed. Information is available via the [Wider Impacts](#) dashboard.

Scottish Ambulance Service (SAS)

Key statistics on unscheduled care operational measures across Scotland, including trends in the number of unscheduled care incidents, responses, conveyances to hospital, response times and hospital turnaround times is available from the [Scottish Ambulance Service \(SAS\)](#) weekly-unscheduled care operational statistics release.

Accident & Emergency (A&E) Activity

Additional information on Accident and Emergency (A&E) performance is available via the weekly [A&E activity and Waiting Times](#) publication, which provides an update of key statistics on attendances at Accident and Emergency (A&E) services across Scotland. Accident and Emergency waiting times and activity reporting on performance against the 4-hour waiting time standard, and the target to reduce attendances at Emergency Departments.

Large decreases in attendances at A&E services in NHS Scotland were observed in spring 2020 winter 2020/21 due to the measures put in place to respond to COVID-19. Since spring, 2021 attendances at A&E have been rising and are getting closer to the pre-COVID levels. However, from the summer of 2021 performance against the four hour standard has dropped below 80% and has remained at this rate for a prolonged period of time.

Emergency Admissions

The information presented in this section aims to provide a better understanding of the underlying trends in emergency admissions during this period.

Figure 17 below shows the overall weekly trend of emergency acute hospital admissions (including COVID-19) from week ending 05 January 2021 to 08 February 2022. The number of emergency admissions have generally been decreasing since week ending 09 November 2021.

Figure 17: Trend of all Emergency Acute Hospital Admissions in Scotland




Table 16 below shows a breakdown of Emergency Admissions to acute hospital across all ages and by age group for the period 15 December 2021 to 08 February 2022.

Table 16: Emergency Hospital Admissions by age as at 08 February 2022³

Age Band	11 January – 18 January	19 January – 25 January	26 January – 01 February	02 February – 08 February
Under 18	1,158	1,283	1,324	1,195
18-29	639	747	745	762
30-39	775	851	818	793
40-49	845	869	860	842
50-54	541	580	587	617
55-59	691	697	697	690
60-64	742	739	780	750
65-69	803	760	826	778
70-74	953	1,013	985	1,002
75-79	912	960	1,018	948
80+	2,144	2,139	2,229	2,115
Total	10,203	10,638	10,869	10,492

Source: RAPID (Rapid and Preliminary Inpatient Data)

3. Please refer to [Appendix 3 – Hospital Admissions Notes](#) for explanatory notes regarding RAPID Hospital Admissions.

In the latest week, there has been a 3.6% decrease in the number of emergency admissions, with those aged 80+ years having the highest number of admissions. Also, in the latest week 53.3% of the hospital admissions related to patients aged 60+.

Waiting Times

Waiting times are important to patients and are a measure of how the NHS is responding to demands for services. Measuring and regular reporting of waiting times highlights where there are delays in the system and enables monitoring of the effectiveness of NHS performance throughout the country.

Public Health Scotland routinely publish a range of statistics on [Waiting Times](#), including: waiting times for diagnostic tests, new outpatient appointments, inpatient and day case treatments.

These statistics continue to be affected by the COVID-19 (Coronavirus) pandemic. At the start of the outbreak, many services were paused or reduced and there were fewer referrals. Boards started to resume relevant services, from June 2020. However, as a second wave of COVID-19 cases emerged through the autumn and winter months, many Boards had to temporarily pause non-urgent diagnostic tests during the months of January and February 2021. Access to services has generally increased since then but some Boards may have been temporarily impacted by a return to high infection rates in recent months as lockdown restrictions eased.

Delayed Discharges

Timely discharge from hospital is an important indicator of quality. It is a marker for person-centred, effective, integrated and harm free care.

For most patients, following completion of health and social care assessments, the necessary care, support and accommodation arrangements are put in place in the community without any delay and the patient is appropriately discharged from hospital.

A delayed discharge occurs when a patient aged 18 years and over, clinically ready for discharge, cannot leave hospital because the other necessary care, support or accommodation for them is not readily accessible and/or funding is not available, for example to purchase a care home place.

Public Health Scotland publish monthly statistics on [Delayed Discharges](#) in Scotland. These figures provide the number of hospital bed days associated with delayed discharges and the number of discharges from hospital following a period of delay. Information is also provided on the number of people experiencing a delay in discharge from hospital at the monthly census point.

Delayed Discharge figures in NHS Scotland have been affected by measures put in place to respond to COVID-19. The marked fall in delayed discharges during 2020 is likely due to patients being moved out of hospital to increase capacity.

Wider Impact of COVID-19

The COVID-19 pandemic has direct impacts on health as a result of illness, hospitalisations and deaths due to COVID-19. However, the pandemic also has wider impacts on health, healthcare, and health inequalities. Reasons for this may include:

- Individuals being reluctant to use health services because they do not want to burden the NHS or are anxious about the risk of infection.
- The health service delaying preventative and non-urgent care such as some screening services and planned surgery.
- Other indirect effects of interventions to control COVID-19, such as changes to employment and income, changes in access to education, social isolation, family violence and abuse, changes in the accessibility and use of food, alcohol, drugs and gambling, or changes in physical activity and transport patterns.

More detailed background information on these potential impacts is provided by the Scottish Public Health Observatory in a section on [Covid-19 wider impacts](#).

The surveillance work stream of the Public Health Scotland social and systems recovery cell aims to provide information and intelligence on the wider impacts of COVID-19 on health, healthcare, and health inequalities that are not directly due to COVID-19. The [wider impact dashboard](#) can be viewed online and includes the following topics:

- Hospital and unscheduled care
- Accident and Emergency attendances
- NHS 24 completed contacts
- Out of hours cases
- Scottish Ambulance Service
- Excess deaths
- Outpatient appointments
- Healthcare for cardiovascular disease
- Healthcare for mental health
- Women booking antenatal care
- Healthcare for birth and babies
- Termination of pregnancy
- Child health
- Cancer
- Substance use
- Injuries

These analyses are based on a selected range of data sources that are available to describe changes in health service use in Scotland during the COVID-19 pandemic. More detailed information is available at NHS Board and Health and Social Care Partnership (HSCP) level.

Contact

Public Health Scotland: p hs.covid19data&analytics@p hs.scot

Further Information

COVID surveillance in Scotland

[Scottish Government](#)

[Daily Dashboard by Public Health Scotland](#) [National Records of Scotland](#)

UK and international COVID reports

[Public Health England](#)

[European Centre for Disease Prevention and Control](#)

[WHO](#)

Weekly National Seasonal Respiratory Report

[Weekly national seasonal respiratory report - Week 5 2022 - Weekly national seasonal respiratory report - Publications - Public Health Scotland](#)

Next Release

The next release of this publication will be 23 February 2022.

Open Data

Data from this publication is available to download from the [Scottish Health and Social Care Open Data Portal](#).

Rate this publication

Let us know what you think about this publication via the link at the bottom of this [publication page](#) on the PHS website.

Early access details

Pre-Release Access

Under terms of the "Pre-Release Access to Official Statistics (Scotland) Order 2008", PHS is obliged to publish information on those receiving Pre-Release Access ("Pre-Release Access" refers to statistics in their final form prior to publication). Shown below are details of those receiving standard Pre-Release Access.

Standard Pre-Release Access:

Scottish Government Health Department
NHS Board Chief Executives
NHS Board Communication leads

Appendices

Appendix 1: Background information

In late December 2019, the People's Republic of China reported an outbreak of pneumonia due to unknown cause in Wuhan City, Hubei Province.

In early January 2020, the cause of the outbreak was identified as a new coronavirus. While early cases were likely infected by an animal source in a 'wet market' in Wuhan, ongoing human-to-human transmission is now occurring.

There are a number of coronaviruses that are transmitted from human-to-human which are not of public health concern. However, COVID-19 can cause respiratory illness of varying severity.

On the 30 January 2020, the World Health Organization [declared that the outbreak constitutes a Public Health Emergency of International Concern](#).

Extensive measures have been implemented across many countries to slow the spread of COVID-19.

Further information for the public on COVID-19 can be found on [NHS Inform](#).

Appendix 2: World Health Organisation (WHO): Contact tracing in the context of COVID-19

The WHO initially produced guidance on "*enhanced criteria to adjust public health and social measures in the context of Covid-19*" in May 2020. The relevant extract from the criteria about the effectiveness of contact tracing within the context of public health surveillance at that time was:

At least 80% of new cases have their close contacts traced and in quarantine within 72 hours of case confirmation	These indicate that the capacity to conduct contact tracing is sufficient for the number of cases and contacts
---	--

Source: <https://apps.who.int/iris/rest/bitstreams/1277773/retrieve>

In response to questions about whether the Scottish Government had been incorrectly comparing Scottish performance with the WHO "standard" (on the basis that counting in Scotland might start at the wrong point in the process), an assessment was undertaken at the start of 2020, and is available within Appendix 2 of the [Weekly Covid-19 Statistical report \(publication date 27 January 2021\)](#).

Please note this "standard" has subsequently been replaced with further [WHO guidance](#) issued in February 2021, reflecting the evolution of the state of the pandemic. This revised guidance now focuses on targeted approaches to contact tracing based on transmission patterns, engaging communities, and prioritising follow-up of high-risk cases when it is not possible to identify, monitor and quarantine all contacts.

Appendix 3: Hospital Admissions Notes

Hospital Admissions

RAPID (Rapid and Preliminary Inpatient Data)

COVID-19 related admissions have been identified as the following: A patient's first positive PCR or LFD (LFD from the 5th of January 2022) for COVID-19 up to 14 days prior to admission to hospital, on the day of their admission or during their stay in hospital. If a patient's first positive test is after their date of discharge from hospital, they are not included in the analysis.

The number reported does not take into account the reason for hospitalisation. Therefore, people that were admitted for a non COVID-19 related reason (and tested positive upon admission) may be included.

RAPID is a daily submission of people who have been admitted and discharged to hospital. These data include admissions to acute hospitals only and do not include psychiatric or maternity/obstetrics specialties. Figures are subject to change as hospital records are updated. It can take 6-8 weeks or longer before a record is finalised, particularly discharge details.

In the data presented here, an admission is defined as a period of stay in a single hospital. There may be multiple admissions for a single patient if they have moved between locations during a continuous inpatient stay (CIS), or if they have been admitted to hospital on separate occasions.

Hospital Inpatients (Scottish Government Data)

Number of patients in hospital with recently confirmed COVID-19, identified by their first positive LFD test (from 5 January 2022) or PCR test. This measure (available from 11 September 2020 and first published 15 September 2020) includes patients who first tested positive in hospital or in the 14 days before admission. Patients stop being included after 28 days in hospital (or 28 days after first testing positive if this is after admission). Further background on this new approach is provided in [this Scottish Government blog](#).

This is based on the number of patients in beds at 8am the day prior to reporting, with the data extract taken at 8am on the day of reporting to allow 24 hours for test results to become available. Where a patient has not yet received a positive test result, they will not be included in this figure. Patients who have been in hospital for more than 28 days and still being treated for COVID-19 will stop being included in this figure after 28 days. All patients in hospital, including in intensive care, and community, mental health and long stay hospitals are included in this figure.

Appendix 4: Contact Tracing

Definitions

An **index case** is generated for each positive result with a test date on or after 28 May 2020. This includes tests derived from Scottish and UK Government laboratories, as well as self-reported LFD's

An **individual** is a unique person who has had a positive test. Before the 5th of January 2022, only positive PCR tests were counted, but LFD tests are included after that point to identify unique individuals. An individual can have multiple positive tests which results in multiple cases within the test and protect system. In these figures, each person is only counted once.

A **contact** may be contacted more than once if multiple positive cases list them as a contact.

Completed cases are cases, which are marked as completed in the case management system, which means that all contacts have been followed up and completed. It excludes cases marked as failed, excluded, in progress or new. In the latest weeks there will be cases which are still open either because contact tracing is still underway (particularly for the latest week) or the NHS Board is still managing the case as part of an open outbreak. Weekly data presented from Monday to Sunday in order to be consistent. Figures are provisional and may change as the test and protect tool is updated by contact tracers.

Individuals unable to be contacted

This information is only available for index cases that have been recorded on the CMS. The CMS went live on 22 June 2020 with NHS Boards migrating on a phased approach with all Boards using CMS from 21 July 2020. Prior to a Board migrating to CMS, data was recorded in a Simple Tracing Tool, which did not give the level of granularity required to report on these measures. These data are developmental and an extensive data quality assurance exercise is underway and data may be revised in subsequent publications. Please note the methodology has changed as of 1 November 2020, a refined method has now been applied to identify unique indexes.

Contact tracers will contact index cases by telephone, and by default, all close contacts will receive an automated SMS. This approach ensures high quality calls can continue to be prioritised for index cases. Even when SMS is defaulted to, in these scenarios, a number of close contacts are still telephoned, following clinical risk assessment, particularly if they are linked to complex cases. When close contacts of index cases are contacted via SMS text message, the GOV.UK Notify Service is used which means it is known if the SMS has been received by the mobile phone, not just that it has been sent. Where the SMS is not received, a contact tracer will attempt to contact the individual through other means.

Appendix 5: Lateral Flow Device Testing

UK Gov other includes any LFD result, which has come through the UK Government route (NHS Digital) which has the test site code "Other". Please note the universal offer results up to 28 July 2021 are reported via this method. From 28 July 2021 onwards, universal offer results are reported separately as Universal Offer.

The Attend An Event, High Cases In Local Area, Lives With Someone Who Is Shielding, Travel Within UK and Universal Offer categories only include data from 28 July 2021 onwards. From this date, these categories are now options when entering a non-work LFD result via the UK Gov portal. Please note that it is up to the user to select the Attend An Event, High Cases In Local Area, Lives With Someone Who Is Shielding or Travel Within UK category, these are not part of any defined testing programme such as Community Testing or University Testing.

University Testing Site tests are tests which took place at a university-testing site, generally in the 2020/21 academic year, though there are still a small number of tests each week in this category. Tests in the university students and university staff categories are tests via the UK

Gov portal for someone entering a test to attend their place of work/education; these tests are from 28th July 2021 onwards and will be for the 2021/22 academic year.

For information regarding LFD testing during term time as part of the Schools Asymptomatic Testing Programme, please visit the [COVID-19 Education Surveillance Report](#).

Please note bulk uploading functionality is not yet available so data is likely to be an undercount. Data will be update and revised in future publications.

Other is any result entered via the [gov.uk website](#) where “none of the above” has been selected. Please note anyone requesting a LFD test via the general population offer, will currently report their results via this category.

Appendix 6: Data sources and Limitations

Date of extraction and analysis

Due to delays in reporting, figures are subject to change as records are updated. A marker (greyed-out block) has been applied where data is preliminary and caution should be taken in their interpretation.

The definitions described below are being used for the purposes of evaluating the impact of the COVID-19 vaccine on COVID-19 cases, COVID-19 related acute hospital admissions and confirmed COVID-19 deaths. The numbers reported in this section use test data, accounting for potential reinfections, and may differ from other sections and elsewhere which only count the number of new COVID-19 cases.

Please note that this section only includes Polymerase chain reaction (PCR) confirmed COVID-19 cases, COVID-19 related acute hospital admissions and confirmed COVID-19 deaths. Lateral flow device (LFD) confirmed cases are not included in this section and therefore will result in an underestimation of recent case rate trends.

COVID-19 PCR test results and associated demographics of an individual are extracted from the Test and Protect database (Corporate Data Warehouse) which contains test results from Electronic Communication of Surveillance in Scotland (ECOSS). Data included in this analysis is reported up until the Friday of the previous week. Non-Scottish residents are excluded from the dataset.

COVID-19 cases are identified as the following: An individual that has tested positive for COVID-19 by PCR. If an individual tests positive more than once, the repeat positive PCR test is only counted if the positive PCR test is more than 90 days apart. Records with missing CHI numbers are excluded as these data cannot be linked to vaccination status.

COVID-19 cases included for the age-standardised rates only includes individuals 10 years old and over. Although the majority of 10 and 11-year-olds are currently not eligible for vaccination, the five-year age band standardised to the 2013 European Standard Population used in this analysis ranges from 10-14 years and therefore cases and denominators for these age groups are included.

Vaccination status for all individuals who test positive for COVID-19 by PCR is extracted from the data used to produce the Public Health Scotland vaccine uptake/daily dashboard. Vaccine records include the number of doses and date of vaccination. Individuals are listed as unvaccinated if there is no vaccination record linked to their unique CHI identifier at the

time of analysis. Vaccination status is taken at date of specimen for COVID-19 cases, acute hospital admissions, or death and assigned to number of doses according to the case definitions described below.

COVID-19 vaccination status is defined as per the following:

- **Dose 1:** An individual that has had one dose of COVID-19 vaccine and has tested positive for COVID-19 by PCR more than 21 days after their first dose of COVID-19 vaccine and have either not had a second dose of COVID-19 vaccine or is less than or equal to 14 days after their second dose of COVID-19 vaccine.
- **Dose 2:** An individual that has had two doses of COVID-19 vaccine and has tested positive for COVID-19 by PCR more than 14 days after their second dose of COVID-19 vaccine and have either not had a booster or third dose of COVID-19 vaccine or is less than or equal to 14 days after their booster or third dose of COVID-19 vaccine.
- **Booster or third dose:** An individual that has had 2 doses of COVID-19 vaccine and a booster or third dose of COVID-19 vaccine and has tested positive for COVID-19 by PCR more than 14 days after their booster or third dose of COVID-19 vaccine.
- **Unvaccinated:** An individual that has had no doses of COVID-19 vaccine and has tested positive for COVID-19 by PCR or has had one dose of COVID-19 vaccine and has tested positive less than or equal to 21 days after their first dose of COVID-19 vaccine.

Denominator population estimates by vaccine status are calculated as the number of people in each vaccine status (defined below) on any given day. For weekly tables, calculations are made for the Friday of each reporting period, whereas for the seven-day rolling average in figures, denominators are calculated daily. Population denominators are calculated as per the definitions below:

- **Dose 1:** The number of individuals that had their first dose of COVID-19 vaccine more than 21 days prior to the reporting period and have either not had a second dose of COVID-19 vaccine or are less than or equal to 14 days after their second dose of COVID-19 vaccine.
- **Dose 2:** The number of individuals that had two doses of COVID-19 vaccine more than 14 days prior to the reporting period and have either not had a booster or third dose of COVID-19 vaccine or are less than or equal to 14 days after their booster or third dose of COVID-19 vaccine.
- **Booster or third dose:** The number of individuals that have had two doses of COVID-19 vaccine and had a booster or third dose of COVID-19 vaccine more than 14 days prior to the reporting period.
- **Unvaccinated:** The unvaccinated population is calculated using the population estimate denominator (all individuals registered with a GP in Scotland) minus the sum of all effective vaccinated individuals in the Dose 1, Dose 2 and third dose/booster populations (described above).

Calculating the denominators for the 16 and over population are taken from the COVID-19 vaccination database. The denominator for under 16-year-olds is from the National Records of Scotland (NRS) mid-2020 population estimates. Population data are extracted from Community Health Index (CHI) dataset representing all those currently registered with a GP

practice in Scotland. These are different denominators than those in the Public Health Scotland COVID-19 Daily Dashboard and may over-estimate the population size as they will include, for example, some individuals who are no longer residents in Scotland. This is a particular issue for the denominator for the unvaccinated cohort, because for vaccinated individuals we know they were resident in Scotland at the time of their vaccination whereas for the unvaccinated cohort there will be a mixture of people who have chosen not to have the vaccine and those who are no longer resident in Scotland. This means that the rates of COVID-19 for the unvaccinated groups will be underestimated, whereas the rates for the vaccinated groups will be more accurate.

COVID-19 related acute hospital admissions have been identified as the following: An individual that has tested positive for COVID-19 by PCR:

- Up to 14 days prior to hospital admission
- On the day of, or day following admission (if no discharge date is available)
- In between hospital admission and discharge (if there is a valid discharge date available)
- Are 15 years old and over.

Where an individual has more than one PCR positive test, positive results are only included for the first PCR positive test associated with a hospitalisation, or if the positive PCR test is more than 90 days after the previous PCR positive test that was eligible for inclusion. Using these criterion, all records of hospitalisation occurring within 90 days of a previous positive test are excluded. Therefore, if a positive PCR test result for an individual meets these criteria for multiple hospital stays, for example, an individual is admitted twice within a week, only the earliest hospital admission is included in the analysis.

If a patient tested positive after their date of discharge from hospital, they are not included in the analysis unless they are readmitted to hospital and meet the criteria described above.

The number of reported acute hospitalisations does not take into account the reason for hospitalisation, Therefore, people that were admitted for a non-COVID-19 related reason (and tested positive upon admission) may be included and result in an overestimation of COVID-19 related acute hospitalisations.

Hospital admission data is extracted from the Rapid and Preliminary Inpatient Data (RAPID) dataset on Monday 07 February 2022. RAPID is a daily submission of people who have been admitted and discharged to hospital. Figures are subject to change as hospital records are updated. Data included in this analysis is reported up until the Friday of the previous week.

In the data presented here, an admission is defined as a period of stay in a single hospital. If the patient has been transferred to another hospital during treatment, each transfer will create a new admission record. Therefore, there may be multiple admissions for a single patient if they have moved between locations during a continuous inpatient stay (CIS), or if they have been admitted to hospital on separate occasions.

Confirmed COVID-19 deaths Death data were extracted from the Scottish Morbidity Records (SMRA) dataset on Thursday 03 February 2022. Data included in these analysis are reported up until the last date of death registration for the previous week.

A confirmed COVID-19 related death is defined as an individual who has tested positive by PCR for SARS-CoV-2 at any time point and has COVID-19 listed as an underlying or

contributory cause of death on the death certificate. Vaccine status is determined at time of most recent specimen date.

Age standardised case, hospitalisation and mortality rates are used to allow comparisons of case, hospitalisation and mortality rates between populations that have different age distributions. The 2013 European Standard Population is used to standardise rates. For more information, see [the Office of National Statistics \(ONS\) methods](#).

COVID Vaccine Wastage:

The single source of vaccination wastage data for Scotland is through an NSS Service Now wastage form, which is populated by health board clinicians which can impact timeliness and accuracy.

It is important to note, that these statistics do not include wastage of vaccines in GPs practices. Therefore, the Scotland level figures reported above may be an under estimate.